

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C07K 14/62, C30B 29/58	A1	(11) International Publication Number: WO 00/01727 (43) International Publication Date: 13 January 2000 (13.01.00)
--	----	--

<p>(21) International Application Number: PCT/DK99/00371</p> <p>(22) International Filing Date: 30 June 1999 (30.06.99)</p> <p>(30) Priority Data: 98610020.4 30 June 1998 (30.06.98) EP 60/092,882 15 July 1998 (15.07.98) US</p> <p>(71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, P.O. Box 3000, DK-2880 Bagsværd (DK).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): MANIQUE, Flemming [DK/DK]; Lerholmvej 39, DK-2750 Ballerup (DK). ILSØE, Christian [DK/DK]; Skovgårds Allé 219, DK-3500 Værløse (DK).</p> <p>(74) Agent: PLOUGMANN, VINGTOFT & PARTNERS A/S; Sankt Annæ Plads 11, P.O. Box 3007, DK-1021 Copenhagen K (DK).</p>	<p>(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW). Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>
---	--

(54) Title: SEEDING CRYSTALS FOR THE PREPARATION OF PEPTIDES OR PROTEINS



(57) Abstract

A method for producing seeding microcrystals for the production of human insulin, said microcrystals being free of non-human pancreatic insulin, the method comprising providing an unseeded suspension of human insulin, said suspension being free of non-human pancreatic insulin, and homogenising said insulin suspension under pressure to result in human insulin microcrystals suitable for use as seeding microcrystals for the production of zinc insulin products. The method of homogenisation under pressure may also be used for the production of seeding microcrystals for other peptides and proteins, in particular pharmaceutical peptides or proteins such as insulin, GLP-1, glucagon and growth hormones.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SEEDING CRYSTALS FOR THE PREPARATION OF PEPTIDES OR PROTEINS

FIELD OF THE INVENTION

- 5 The present invention relates to seeding crystals for the preparation of peptides or proteins such as zinc insulin products.

BACKGROUND OF THE INVENTION

- 10 The "Lente" family of zinc insulin products are insulin zinc suspensions of the type originally developed in the 1950's with the aim of producing insulin preparations that would be able to cover diabetics' insulin requirement with a single daily injection ((see e.g. Jens Brange, *Galenics of Insulin*, 1987). Various Lente insulin products having different action profiles are available in the form of different combinations of amorphous and/or crystalline insulin
- 15 particles from Novo Nordisk A/S, Denmark. These include SEMILENTE, a suspension of amorphous insulin particles, ULTRALENTE, a suspension of crystalline insulin particles, and LENTE, which is a mixture of 30% amorphous and 70% crystalline insulin particles.

- For several decades, seeding crystals for preparation of the "Lente" zinc insulin products
- 20 have been prepared by the same basic freeze-drying method that was developed and patented in the early 1950's. This method, which is described in GB patent specification No. 766,994, involves the addition of freeze-dried amorphous insulin, typically beef insulin, to an insulin-containing crystallisation medium to result in the formation of a suspension of insulin microcrystals of a size of about 2-7 μm . This suspension, which is eventually used
- 25 for the preparation of the final crystalline zinc insulin product, is filled into small vials (e.g. 10 ml), frozen in an alcohol/carbon dioxide mixture and stored frozen at a temperature at or below -18°C .

Although still in use, this method has a number of disadvantages:

30

1. It is based on the use of beef insulin, since it has until now not been possible to produce acceptable microcrystals of pure human insulin. As a result of the use of beef insulin nuclei for the formation of the microcrystals, the end product contains a small amount of beef insulin, which is undesirable.

35

2. The freeze-drying method requires a lyophiliser and subsequent shipping and storage at a temperature of no more than -18°C . This is expensive and requires a great deal of space.

3. The method of preparation is extremely difficult to perform in a sufficiently aseptic
5 manner.

It would therefore be advantageous to be able to produce insulin seeding crystals using a method which does not suffer from the disadvantages of the known methods. It has now surprisingly been found that it is possible, using a relatively simple and inexpensive
10 process, to produce insulin seeding crystals which are free of beef insulin, which can be stored at room temperature and which result in insulin preparations having advantageous properties in terms of e.g. crystal particle size and uniformity. Furthermore, it is also contemplated that this process will be applicable to the production of seeding crystals for other peptides and proteins, in particular peptides or proteins used as pharmaceuticals.

15

SUMMARY OF THE INVENTION

It is thus an object of the present invention to provide a novel method for the production of peptide or protein seeding crystals. More particularly, it is an object of the invention to
20 provide a method for the production of insulin seeding crystals which does not require the use of beef insulin, which makes possible storage and transport without the need for expensive freeze-drying and storage at sub-zero temperatures, and which can be performed in a closed system so as to more readily allow the use of aseptic production methods.

25

Another object of the invention is to provide a method for the preparation of insulin seeding crystals for the production of crystalline zinc insulin suspensions having a narrow particle size distribution.

30 In its broadest aspect, the present invention thus relates to a method for producing seeding microcrystals for the production of a peptide or protein, comprising providing an unseeded suspension of a peptide or protein and homogenising said suspension under pressure to result in peptide or protein microcrystals suitable for use as seeding microcrystals.

In a particular embodiment, the invention relates to a method for producing seeding microcrystals for the production of human insulin, said microcrystals being free of non-human pancreatic insulin, comprising providing an unseeded suspension of human insulin, said suspension being free of non-human pancreatic insulin, and homogenising said insulin suspension under pressure to result in human insulin microcrystals suitable for use as seeding microcrystals for the production of zinc insulin products.

Another aspect of the invention relates to a method for the production of a peptide or protein product, comprising providing an unseeded suspension of a peptide or protein and seeding said suspension with microcrystals produced by the method indicated above.

In a particular embodiment of this aspect of the invention, the peptide or protein product to be produced is a zinc insulin product, and the unseeded suspension is a suspension of human insulin.

15

A further aspect of the invention relates to human insulin microcrystals suitable for use as seeding microcrystals for the production of zinc insulin products, said microcrystals being free of non-human pancreatic insulin.

20 A still further aspect of the invention relates to human zinc insulin product free of non-human pancreatic insulin.

DETAILED DESCRIPTION OF THE INVENTION

25 As indicated above, the method of the invention is directed to the production of seeding microcrystals for peptides and proteins in general, in particular for peptides and proteins that are used as pharmaceuticals. More particularly, the method is directed to seeding microcrystals for the production of therapeutic peptides or proteins such as insulin, GLP-1, glucagon, and growth hormones such as human growth hormone, as well as analogues and derivatives of such peptides and proteins. The peptide or protein is in particular human insulin or an analogue or derivative thereof as described below. In the context of peptides and proteins other than insulin, the terms "analogue" and "derivative" are to be understood analogously to the definitions given below in the context of insulin.

As used in the present text, the term "human insulin" is used to designate naturally occurring human insulin as well as insulin analogues and insulin derivatives. The term "insulin analogue" is used to designate a peptide with insulin activity, derived from a naturally occurring insulin by substitution of one or more amino acid residues, deletion of
5 one or more amino acid residues and/or addition of one or more amino acid residues. An insulin or insulin analogue may optionally be in the form of an "insulin derivative", the term "derivative" referring to a peptide in which one or more of the amino acid residues of the parent peptide have been chemically modified, e.g. by alkylation, acylation, ester formation or amide formation. An "acylated insulin" (or insulin analogue) is an insulin (or insulin
10 analogue) which has an acyl group in the ϵ -amino group of one or more amino acid residues, often a lysine residue.

As used herein, the term "non-human pancreatic insulin" refers to naturally occurring insulin from a non-human source, e.g. bovine or porcine insulin.

15

The basic principle of a presently preferred embodiment of the invention is shown schematically in Fig. 1. The apparatus of Fig. 1 comprises a receiving vessel 1, from which the insulin suspension is transferred by means of a pump 2 into a homogeniser 3. The homogeniser 3 comprises a valve with a very small opening through which the insulin
20 suspension is pumped at a high pressure, e.g. about 1000 bars or higher. Upon exiting the valve, the insulin suspension is subjected to a sudden drop in pressure, which results in the rupture of the insulin crystals, i.e. a homogenisation effect. Since the insulin suspension is preferably subjected to multiple homogenisation cycles in order to result in a sufficiently homogenous suspension of microcrystals having the desired particle size and size
25 distribution, and since the high pressure used in the homogeniser 3 results in an increase in the temperature of the suspension, the apparatus preferably also comprises a heat exchanger 4 downstream of the homogeniser 3 in order to reduce the suspension temperature. From the heat exchanger 4, the insulin suspension is returned to the receiving vessel 1 for further homogenisation cycles as necessary.

30

The temperature of the suspension increases according to the following equation:

$$\Delta T = P/(c \times \delta)$$

where:

5

ΔT = temperature increase ($^{\circ}\text{C}$)

P = suspension pressure ($\text{N} \times \text{m}^{-2}$)

δ = suspension density ($\text{g} \times \text{m}^{-3}$)

c = specific heat ($\text{J} \times \text{g}^{-1} \times ^{\circ}\text{C}^{-1}$)

5

The pressure and temperature are monitored during the process, and the above equation can be used in connection with design of the apparatus and regulation of the process.

In the method according to the invention, homogenisation is typically performed at a pressure of at least about 500 bars, preferably at least about 800 bars, more preferably at least about 1000 bars. In certain cases, the pressure may e.g. be at least about 1200 bars, for example up to about 1500 bars or more, even though such high pressures of above about 1000 bars are generally not believed to be necessary.

15 In a preferred embodiment, homogenisation of the suspension is performed using multiple homogenisation cycles, i.e. at least 2 cycles, since the use of multiple homogenisation cycles has been found to provide improved results, i.e. optimisation of seeding crystal size and uniformity. It is thus contemplated that it will normally be advantageous to use more than 2 cycles, such as 3, 4, 5, 6, 7, 8, 9 or 10 cycles or even more, e.g. in certain cases up
20 to 15 or 20 cycles or perhaps even more than 20 cycles. The most advantageous number of homogenisation cycles will be determined by the person skilled in the art in each individual case based on factors such as the nature of the insulin suspension, the nature of the homogenisation apparatus used, the pressure used for homogenisation, and the desired insulin microcrystal particle size and size distribution.

25

Since, as indicated above, the high pressure used in the homogeniser results in an increase in the temperature of the suspension, the use of multiple homogenisation cycles is preferably accompanied by the use of a heat exchanger in order to reduce the suspension temperature, so that the suspension is maintained at a suitable temperature throughout the
30 homogenisation process. Such heat exchangers are known in the art, and the person skilled in the art will readily be able to adapt the characteristics of the heat exchanger to suit the given process and apparatus. Preferably, the temperature of the recycled insulin suspension is maintained within the range of about 10–40 $^{\circ}\text{C}$, e.g. about 20–35 $^{\circ}\text{C}$.

Although the particle size of the resulting insulin microcrystals will vary depending on the intended use, suitable microcrystals will often have an average particle size, as defined by the longest diagonal of the crystals, in the range of about 0.5-4 μm , e.g. about 1-3 μm .

5 The result of the homogenisation process is human insulin microcrystals suitable for use as seeding microcrystals for the production of zinc insulin products, the microcrystals having the important feature of being free of non-human pancreatic insulin. For the production of zinc insulin products, the seeding microcrystals of the invention will be used in a conventional manner, i.e. an unseeded suspension of human insulin is seeded with the
10 suspension of microcrystals produced as described above, and crystallisation is allowed to proceed in a manner known *per se* in the art. As is normal in the art, the precise amount of microcrystals to be added to a given unseeded insulin suspension may be determined empirically.

15 The invention will be further illustrated by the following non-limiting examples.

EXAMPLES

Materials and methods

20

Using the basic homogenisation process and apparatus described above, i.e. a recirculating homogeniser equipped with a heat exchanger, a number of experiments were performed to test the effect of the number of homogenisation cycles as well as homogenisation pressure and time.

25

The apparatus used was a Rannie high pressure homogeniser, model LAB 10,51 VH (series 1.89239), equipped with a ceramic valve, type SEO 719685. The capacity of the homogeniser was 80 l/h at a pressure of 1000 bars. A centrifugal pump provided an inlet pressure of 4.5-5 bars. The heat exchanger for these experiments used a cooling water
30 temperature of about 20°C. However, since the capacity of the heat exchanger was insufficient in relation to this particular homogeniser, the outlet temperature of the insulin suspension was somewhat higher at the maximum homogeniser output, i.e. about 28-29°C, but slightly lower at a lower homogeniser output of about 65 l/h, i.e. about 24-28°C. The receiving vessel comprised a 100 l pressure tank and a small conic vessel with a volume of
35 about 3 l.

The insulin suspension used for producing the microcrystals was a pooled batch (2 x 20 l) of unseeded ULTRALENTE HM(ge), 100 U/ml, from Novo Nordisk A/S.

5 EXAMPLE 1

A 10 l portion of the pooled batch of the ULTRALENTE insulin suspension was homogenised at a pressure of 1000 bars, the suspension being recirculated for multiple homogenisation cycles as described above, resulting in a gradually increased degree of homogenisation. A flow rate of 80 l/h was used. A total of 18 homogenisation cycles were performed, and samples were taken for the first 10 cycles and after the final cycle. The temperature of the insulin suspension was measured in the outlet conduit between the homogeniser and the heat exchanger. The times and measured temperatures for the various cycles were as indicated in Table 1 below:

15

Table 1

Example number	Number of homogenisation cycles	Time from start (minutes)	Temperature (°C)
1-0	0	0	12.5
1-1	1	5	29
1-2	2	12	29
1-3	3	19	28.3
1-4	4	26	29.5
1-5	5	33	29.5
1-6	6	40	29.8
1-7	7	47	29.5
1-8	8	54	29.2
1-9	9	61	28.1
1-18	18	122	28.1

A number of the samples were investigated by microscope, and the following observations were made:

20

Example 1-0: mostly whole and sharp-edged rhombohedral crystals having a size of about 3-80 μm ; some broken crystals and crystal fragments.

Example 1-1: still many whole rhombohedral crystals having a size of about 20-40 μm , but
5 also many small crystal fragments with a size of 3 μm or less.

Example 1-2: still some whole rhombohedral crystals with a size of up to about 20 μm as well as a few larger crystal agglomerations of up to about 40 μm ; even more small crystal fragments of 3 μm or less.

10

Example 1-18: small microcrystals of about 1 μm or less; a few crystal fragments of up to about 10 μm ; no whole rhombohedral crystals.

EXAMPLE 2

15

A 5 l portion of the pooled batch of the ULTRALENTE insulin suspension was homogenised at a pressure of 1000 bars, the suspension being recirculated for multiple homogenisation cycles as described above, using a flow rate of 65 l/h. A total of 10 homogenisation cycles were performed, and samples were taken after each cycle. In this
20 case, instead of being led directly back to the receiving vessel from the heat exchanger, the suspension was collected after each cycle. A sample was taken from each portion, and the remainder of the portion was returned to the receiving vessel for the next homogenisation cycle. The times and measured temperatures were as follows:

Table 2

Example number	Number of homogenisation cycles	Time from start (minutes)	Temperature (°C)
2-0	0	0	-
2-1	1	-	24.5
2-2	2	-	26.3
2-3	3	-	27.1
2-4	4	15	27.7
2-5	5	-	27.8
2-6	6	-	28.0
2-7	7	-	28.0
2-8	8	-	28.0
2-9	9	-	-
2-10	10	30	-

EXAMPLE 3

5 In order to investigate the effect of the homogenisation pressure, tests were performed at 1400-1500 bars, with a total of 9 homogenisation cycles. Due to the increased pressure and the accompanying increased temperature of the suspension, the flow rate was further reduced to 54 l/h to allow the heat exchanger to provide a sufficiently reduced temperature. The batch size in this case was 3 l. The temperature of the suspension was maintained at 10 about 26-29°C.

Table 3

Example number	Number of homogenisation cycles	Time from start (minutes)	Temperature (°C)
3-0	0	0	16.3
3-9	9	30	28.6

EXAMPLE 4

The same procedure as in Example 3 was used, with the exception that the ULTRALENTE insulin suspension in this case had formed divergent crystals ("roses") during crystallisation. The batch size was 2 l, and the homogenisation time was therefore reduced correspondingly to a total of 21 minutes.

Table 4

Example number	Number of homogenisation cycles	Time from start (minutes)	Temperature (°C)
4-0	0	0	-
4-9	9	21	27.7

10

EXAMPLE 5*Seeding experiments with selected batches of microcrystals*

15 Seeding experiments were performed to test selected batches of the human insulin microcrystals prepared as described above. As a reference, a standard bovine microcrystal seeding batch was also tested. These experiments were performed using 1 l batches of ULTRALENTE (40 U/ml). Crystallisation was performed using propeller agitation for a period of 20 hours. The results are shown in Table 5 below.

20

Table 5

Microcrystals of Example number	Average crystal size (µm)	10%-90% Deviation (µm)
Reference	28	16
1-18	23	15
2-10	26	17
3-9	26	17
4-9	27	16

It may be seen from the results in Table 5 that the human insulin microcrystals prepared according to the invention gave an insulin crystal size and deviation comparable to that obtained using the standard bovine microcrystals, and that the five seeding batches gave
5 largely identical results.

The five insulin batches prepared as described above were in addition analysed with regard to a number of other parameters, including pH, insulin strength, A+M+B component, percentage of amorphous insulin, content of methyl para-hydroxybenzoate, dimer and
10 polymer content, acidic and neutral desamidoins content, and zinc content. It was found that insulin batches prepared using microcrystals according to the invention were generally comparable to insulin prepared using the standard bovine microcrystals.

Since it is known that the crystallisation time and type of agitation can have an effect on the
15 appearance of the rhombohedrons that are formed, a single batch prepared according to the invention (Example 3-9) was used for seeding tests in which the crystallisation time and type of agitation were varied. With regard to agitation, no substantial differences were observed between crystals obtained using propeller agitation and agitation using "cradle
movements". With regard to crystallisation time, it was found that 4 hours was sufficient,
20 i.e. a crystallisation time of 20 hours was found to be unnecessary. There was a tendency for the best results to be obtained using propeller agitation and a crystallisation time of 4 hours, as this led the least amount of deviating crystals.

Conclusion

25

It may be concluded that pure microcrystals of human insulin can be produced by high pressure homogenisation of an unseeded ULTRALENTE HM(ge) preparation. This method results in microcrystals in the form of small crystal fragments with a particle size of about 1-
2 μm and some larger fragments with a particle size of up to about 10 μm .

30

Varying the pressure from 1000 bars to about 1500 bars did not have any noticeable effect on the microcrystals. On the other hand, the number of homogenisation cycles has an effect, at least up to a point, an increased number of cycles resulting in a more uniform
microcrystal suspension with a larger proportion of microcrystals having a size of about 1-2
35 μm and a smaller proportion of larger crystal fragments and whole rhombohedric crystals.

However, the number of homogenisation cycles required to result in a given degree of homogenisation is also related to the homogenisation time per cycle.

The increase in temperature of the suspension measured in these experiments as a result
5 of the homogenisation process did not appear to effect the microcrystals in terms of
chemical degradation. Temperature regulation can be optimised by suitable changes in
e.g. the design and capacity of the heat exchanger.

The variation in the particle size of the microcrystals produced in these tests, and thus the
10 variation in particle size of the zinc insulin product prepared using the microcrystals, could,
if desired or necessary, be reduced by means of e.g. sedimentation or centrifugation.

The seeding qualities of the microcrystals produced according to the invention have been
shown to be acceptable, since the microcrystals result in zinc insulin products with
15 rhombohedral crystals having an acceptable crystal size.

CLAIMS

1. A method for producing seeding microcrystals for the production of a peptide or protein, comprising providing an unseeded suspension of a peptide or protein and homogenising
5 said suspension under pressure to result in peptide or protein microcrystals suitable for use as seeding microcrystals for the production of said peptide or protein.
2. The method of claim 1, wherein the peptide or protein is a pharmaceutical peptide or protein.
10
3. The method of claim 2, wherein the peptide or protein is selected from the group consisting of insulin, GLP-1, glucagon, and growth hormones such as human growth hormone, as well as analogues and derivatives thereof.
- 15 4. The method of claim 1 for producing seeding microcrystals for the production of human insulin, said microcrystals being free of non-human pancreatic insulin, comprising providing an unseeded suspension of human insulin, said suspension being free of non-human pancreatic insulin, and homogenising said insulin suspension under pressure to result in human insulin microcrystals suitable for use as seeding microcrystals for the production of
20 zinc insulin products.
5. The method of any of claims 1-4, wherein homogenisation is performed at a pressure of at least about 500 bars, preferably at least about 800 bars, more preferably at least about 1000 bars, e.g. at least about 1200 bars.
25
6. The method of any of claims 1-5, wherein homogenisation is performed using multiple homogenisation cycles.
7. The method of claim 6, wherein the temperature of the recycled suspension is controlled
30 using a heat exchanger.
8. The method of claim 7, wherein the temperature of the recycled suspension is maintained within the range of about 10-40°C, e.g. about 20-35°C.

9. The method of any of claims 1-8, wherein the microcrystals have an average particle size, as defined by the longest diagonal of the crystals, in the range of about 0.5-4 μm , e.g. about 1-3 μm .
- 5 10. Human insulin microcrystals produced according to the method of any of claims 4-9.
11. Human insulin microcrystals suitable for use as seeding microcrystals for the production of zinc insulin products, said microcrystals being free of non-human pancreatic insulin.
- 10 12. Microcrystals according to claim 11 having an average particle size in the range of about 0.5-4 μm , e.g. about 1-3 μm .
13. A method for the production of a peptide or protein product, comprising providing an
15 unseeded suspension of a peptide or protein and seeding said suspension with microcrystals produced by the method of any of claims 1-9.
14. The method of claim 13, wherein the peptide or protein product to be produced is a zinc
20 insulin product, the unseeded suspension is a suspension of human insulin, and the seeding microcrystals are human insulin microcrystals.
15. A human insulin product free of non-human pancreatic insulin, produced by the method of claim 14.
- 25 16. A human zinc insulin product free of non-human pancreatic insulin.




Fig. 1

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/DK 99/00371

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07K14/62 C30B29/58

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07K C30B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 096 631 A (NOELLE ANCENIS COOP AGRICOLE) 21 December 1983 (1983-12-21)	
A	EP 0 582 351 A (HOLLAND SWEETENER CO) 9 February 1994 (1994-02-09)	
A	PATENT ABSTRACTS OF JAPAN vol. 017, no. 100 (C-1030), 26 February 1993 (1993-02-26) & JP 04 288013 A (KYORIN PHARMACEUT CO LTD), 13 October 1992 (1992-10-13) abstract	
	-/-	

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "Z" document member of the same patent family

Date of the actual completion of the international search

15 October 1999

Date of mailing of the international search report

22/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo rt,
Fax: (+31-70) 340-3018

Authorized officer

Cervigni, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 99/00371

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DEMATTEI R C ET AL: "CONTROLLING NUCLEATION IN PROTEIN SOLUTIONS" JOURNAL OF CRYSTAL GROWTH, vol. 122, no. 1 / 04, 2 August 1992 (1992-08-02), pages 21-30, XP000306488 ISSN: 0022-0248	
X	GB 766 994 A (NOVO TERAPEUTISK LABORATORIUM) 30 January 1957 (1957-01-30) the whole document page 2, column 2, line 65 - line 101; examples	10-12, 15,16
X	GB 766 995 A (NOVO TERAPEUTISK LABORATORIUM) 30 January 1997 (1997-01-30) the whole document examples	10-12, 15,16
X	WO 88 02633 A (NOVO INDUSTRI AS) 21 April 1988 (1988-04-21) the whole document	10-12, 15,16
X	WO 90 00176 A (BIOBRAS BIOQUIMICA DO BRASIL S) 11 January 1990 (1990-01-11) abstract	10-12, 15,16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 99/00371

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0096631	A	21-12-1983	FR 2527903	A 09-12-1983
			AT 21322	T 15-08-1986
			CS 9102258	A 15-07-1992
			EG 16929	A 30-03-1989
			ES 522907	A 01-08-1984
			OA 7453	A 31-12-1984
			PT 76781	A,B 01-06-1983
			SK 278462	B 04-06-1997
EP 0582351	A	09-02-1994	NL 9201408	A 01-03-1994
			AT 175208	T 15-01-1999
			CN 1092429	A,B 21-09-1994
			DE 69322820	D 11-02-1999
			DE 69322820	T 19-08-1999
			MX 9304719	A 31-05-1994
			US 5502238	A 26-03-1996
JP 04288013	A	13-10-1992	JP 2554784	B 13-11-1996
GB 766994	A		NONE	
GB 766995	A		NONE	
WO 8802633	A	21-04-1988	AU 8175387	A 06-05-1988
			DK 332188	A 17-06-1988
			EP 0265214	A 27-04-1988
			FI 882921	A 17-06-1988
			JP 1501389	T 18-05-1989
			PT 85950	A 01-11-1987
			ZA 8707826	A 21-04-1988
WO 9000176	A	11-01-1990	NONE	